



10-19	50mg every second day
< 10	50 mg once weekly

The grade of renal impairment and / or the serum creatinine concentration may be approximately equated to creatinine clearance levels as indicated below.

Creatinine clearance ml/min /1.73 m <sup>2</sup>	Approx. * serum creatinine (micromole/l)	Renal impairment Grade
> 60	-	-
20 to 59	150 to 300	Mild
10 to 19	300 to 700	Moderate
< 10	> 700	Severe

\* Serum creatinine concentration is only an approximate guide to renal function unless corrected for age, weight and sex.

#### **CONTRAINDICATIONS:**

Proguanil should be used with caution in patients with severe renal impairment.

#### **WARNING AND PRECAUTIONS:**

Proguanil should be used with caution in patients with several renal impairments. There have been rare reports of haematological changes in such patients. In any locality where drug resistant malaria is known or suspected. It is essential to take local medical advice on what prophylactic regimen is appropriate. Prophylactic use of Proguanil alone may not be sufficient.

#### **DRUG INTERACTIONS:**

Antacids may reduce the absorption of proguanil, so should be taken at least 2-3 hours apart.

#### **SIDE EFFECTS:**

Side effect which may occur in adult patients include, but are not limited to abdominal pain, nausea, vomiting, headache, diarrhea, asthenia, dizziness, anorexia.

#### **KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT:**

The following effects have been reported in cases of over dosage, haematuria, epigastric discomfort and vomiting. There is no specific antidote and symptoms should be treated as they arise.

#### **STORAGE:**

Store in a cool and dry place.

Protect from heat and light.

**KEEP MEDICINE OUT OF REACH OF CHILDREN.**

**Presentations:**

Available in Blister of 10 tablets in the packing of 10x10 tablets carton.